

# Mapping and Prediction of Coal Workers' Pneumoconiosis with Bioavailable Iron Content in the Bituminous Coals

Xi Huang, Weihong Li, Michael D. Attfield, Arthur Nádas,  
Krystyna Frenkel, and Robert B. Finkelman

doi:10.1289/ehp.7679 (available at <http://dx.doi.org/>)  
Online 15 April 2005



## **Mapping and Prediction of Coal Workers' Pneumoconiosis with Bioavailable Iron Content in the Bituminous Coals**

Xi Huang<sup>1\*</sup>, Weihong Li<sup>1</sup>, Michael D. Attfield<sup>2</sup>, Arthur Nádas<sup>1</sup>, Krystyna Frenkel<sup>1</sup>, and Robert B. Finkelman<sup>3</sup>

<sup>1</sup>Department of Environmental Medicine and NYU Cancer Institute, New York University School of Medicine, 550 First Avenue, PHL Room 802, New York, NY 10016; <sup>2</sup>Division of Respiratory Diseases Studies, National Institute for Occupational Safety and Health, Center for Disease Control and Prevention, Morgantown, WV 26505; <sup>3</sup>US Geological Survey, M.S. 956 National Center, 12201 Sunrise Valley Drive, Reston, VA 20192

\*. To whom correspondence should be addressed.

Tel. (212)-263-6650

Fax: (212)-263-6649

Email: [xihuang@env.med.nyu.edu](mailto:xihuang@env.med.nyu.edu)

**Running Title:** Mapping and Predicting Coal's Toxicity

**Keywords:** Bioavailable iron, pneumoconiosis, coal, calcite, COPD

**Acknowledgement:** This project was supported by grant OH 03561 from the National Institute for Occupational Safety and Health (NIOSH) and in part by the National Institute of Environmental Health and Sciences Center Grant ES00260.

**Abbreviations:** BAI: Bioavailable iron; COPD: Chronic obstructive pulmonary disease; CWP: Coal workers' pneumoconiosis; USGS: US Geological Survey; NSCWP: National Study of Coal Workers' Pneumoconiosis.

## Outline of section headers:

Abstract -----	4
Introduction -----	5 - 6
Materials and Methods -----	6 - 9
Results -----	9 - 11
Discussion -----	11 - 14
References -----	15 - 19
Figure Legends -----	20
Tables -----	21 - 23
Figures -----	24 - 25

**Abstract:**

Based on the first National Study of Coal Workers' Pneumoconiosis (CWP) and the US Geological Survey database of coal quality, we show that the prevalence of CWP in seven coalmine regions correlates with levels of bioavailable iron (BAI) in the coals from that particular region [correlation coefficient  $r=0.94$ ,  $p<0.0015$ ]. CWP prevalence is also correlated with contents of pyritic sulfur ( $r=0.91$ ,  $p<0.0048$ ) or total iron ( $r=0.85$ ,  $p<0.016$ ), but not with coal rank ( $r=0.59$ ,  $p<0.16$ ) or silica ( $r=0.28$ ,  $p<0.54$ ). BAI was calculated using our model taking into account chemical interactions of pyrite, sulfuric acid, calcite, and total iron. That is, iron present in coals can become bioavailable by pyrite oxidation, which produces ferrous sulfate and sulfuric acid. Calcite is the major component in coals that neutralizes the available acid and inhibits iron's bioavailability. Therefore, levels of BAI in the coals are determined by the available amounts of acid after neutralization of calcite and the amount of total iron in the coals. Using the linear fit of CWP prevalence and the calculated BAI in the seven coalmine regions, we have derived and mapped seven thousand coal samples' pneumoconiotic potencies. Our studies indicate that levels of BAI in the coals may be used for the prediction of coal's toxicity, even before large-scale mining.